8092-EN-01 DTIC.

CHARACTERISATION OF BACKGROUND BIOLOGICAL AEROSOL

S.G. JENNINGS

(Principal Investigator) University College Galway

CONTRACT NUMBER: N68171-96-C-9124

3rd Interim Report

April 1997 - June 1997

Approved for public releases

Distribution Unlimited

DTIC QUALITY INSPECTED 4

The research reported in this document has been made possible through the support, and sponsorship of the U.S. Government through its European Research Office of the U.S. Army. This report is intended only for the internal management use of the Contractor and the U.S. Government.

19970911 091

REPORT DOCUMENTATION PAGE OMB NO 0704-0188 Figure legaling burger by this joileding of information is climated to account to the composition of information of information in composition of information and information of informati Constitutes forth 104 Association as 11101-101. July 30 1997 J. REPORT TYPE AND DATES COVERED Interim Report: Apr'97-Jun'97 1. AGENCY USE ONLY (Leave blank) J. FUNDING NUMBERS 4. TITLE, AND SUBTITLE Characterisation of Background N68171-96-C-9124 Biological Aerosol 6. AUTHORIS) S. G. Jennings and C. M. Kenny 8. PERFORMING ORGANIZATION 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 0003 University College Galway, Ireland. 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING , MONITORING AGENCY REPORT NUMBER U.S. Army Research, Development and Standardisation Group, 223 Old Marylebone Road, London NW1 5TH, U.K. 11. SUPPLEMENTARY NOTES None 125 DISTRIBUTION CODE 123. DISTRIBUTION / AVAILABILITY STATEMENT Unlimited 13. ABSTRACT (Maximum 200 words) Sampling of ambient air using a glass cyclone system for fluorescence background determination is described. Weekly samples over 12 hour sampling periods are taken at the University College Galway's atmospheric research field station at Mace Head, on the west coast of Ireland. The bioaerosol sampling system and procedures used have previously been discussed in the 1st Interim Report. Use is made of a fluorescence protocol developed in the ERDEC Laboratories. Representative fluorescence excitation/emission spectra are presented and compared for both background and polluted conditions. Intercomparison between Winter and Spring signatures is made. 15. NUMBER OF PAGES 14. SUBJECT TERMS 16 16 PRICE CODE SECURITY CLASSIFICATION OF ABSTRACT 20. LIMITATION OF ABSTRACT SECURITY CLASSIFICATION 17. SECURITY CLASSIFICATION OF THIS PAGE OF REPORT

Unclassified

Unclassified

Unclassified NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89) Protection by ANN No. 239-18 199-101

None

Form Approved

CHARACTERISATION OF BACKGROUND BIOLOGICAL AEROSOL

Abstract

Bioaerosol sampling has been been carried out at Mace Head on the West coast of Ireland on a weekly basis. Sampling time has been increased from a three hour to a twelve hour period as from January 31st, 1997. Representative fluorescence excitation/emission spectra are presented for both background and polluted air conditions over the period from 11/29/96 to 05/08/97.

1. Introduction

Biological monitoring is carried out at Mace Head for three reasons, firstly to determine the influence of the Atlantic ocean on bioactivity, secondly to establish a "baseline" for future remote biological sensing systems and thirdly for the investigation of the correlation of bioaerosol data with non-viable data already available at Mace Head. The method currently in use at Mace Head for capturing airborne biological material is an impinger system and the model type is an Aerojet General glass cyclone. The impinger system relies on the fact that a known quantity of air is pulled at a high velocity through a fluid in a glass vessel. Aerosolised biological matter are retained in the washing fluid while the air passes through the fluid and out through the pump system. The high, almost sonic air speed used in impinger devices results in high shear forces which breakup bacterial/particulate aggregates so that the total count obtained closely reflects the actual number of viable organisms. This shearing action differentiates impingement from other methods, all of which measure only the number of bacteria bearing spores which give colony forming units (cfu). Sampling and fluorescence measurement protocols used have been documented in the 1st Interim report.

2. Background Aerosol Characterisation

Twelve hourly bioaerosol samples were collected at Mace Head from 01/31/97 onwards, replacing the previous 3 hour sampling period. Sampling times in Greenwich Mean Time (GMT) are given in Table 1. Representative emission spectra based on the Winter months of November, December, January and February and the Spring months of March, April and May are shown in Figure 1 through Figure 3. Data includes bioaerosol samples from

both marine (wind sector 180-300°) and continental (wind sector 45-135°) air masses, and are summarised in Table 2. In addition to the condition that the wind direction is between 180 and 300°, a further constraint was imposed to ensure marine air mass conditions. It was required that the condensation nuclei (CN) number concentration be < 700 cm⁻³.

Table 1. Bioaerosol sampling times.

Date	Sampling Times-GMT
11/29/96	11:15-14:15
12/06/96	11:15-14:15
01/09/97	11:45-14:45
01/13/97	10:30-13:30
01/31/97	01:00-13:00
02/07/97	01:00-13:00
03/13/97	04:00-16:00
03/19/97	21:00-09:00
04/03/97	04:00-16:00
04/24/97	04:00-16:00
05/02/97	05:00-17:00
05/08/97	04:00-16:00

Table 2. Bioaerosol measurement periods.

Season	Date	Wind Sector
Winter	11/29/96	Marine
	12/06/96ª	Modified marine
	01/09/97	Continental
	01/13/97	Marine
	01/31/97	Continental
	02/07/97 ^b	Modified marine
Spring	03/13/97	Marine
	03/19/97	Marine
	04/03/97 ^c	Modified marine
	04/24/97	Marine
	05/02/97 ^d	Modified continental
	05/08/97°	Modified marine

a: Wind direction was 150° for the 3 hour period.

b: Wind direction was 300-324° for 3 hours.

c: Wind direction was 300-312° for 9 hours.

d: Wind direction was 150° for 1 hour.

e: Wind direction was 300-310° for 7 hours.

Emission spectra have been grouped by the excitation wavelengths 282, 350 and 450 nm. Back trajectory data for all selected dates are displayed in Figure 4 (a) to Figure 4 (l), courtesy of Peter Lynch and Liam Campbell at Met Éireann. The calculations for each trajectory are based on a 4 day period up to midday of the specified dates.

Marine and modified marine fluorescence intensity values at excitation wavelength 282 nm do not show any significant increase in relation to increased sampling time over the Winter-Spring period. An exception occurs on the 05/08/97 (Figure 1 (k)) where there is a comparitively large increase in intensity during a modified marine sampling period. Hourly CN number concentration for the same period were < 700 cm⁻³ and do not show any dominant continental influences.

The highest continental signal at 282 nm occurs on the 01/09/97, (Figure 1 (d)). Continental fluorescence intensity values are relatively low on the 01/31/97 (Figure 1 (f)) and 05/02/97 (Figure 1 (l)). Intensity values on the 01/31/97 are lower than the adjacent modified marine (Figure 1 (e)) sampling period.

At 350 nm the fluorescence emission signal for the continental air masses - Figure 2 (d), Figure 2 (f) and Figure 2 (l) exceed that of the marine signal (Figure 2 (c)) and that of modified marine signals (Figure 2 (e), Figure 2 (k)) in comparing measurements taken at fairly close periods to one another. The greatest fluorescence occurs on 01/09/97 for a continental air mass, similar to that for excitation at 282 nm.

Fluorescence intensity values at 450 nm are extremely low with the strongest signal occurring on the 01/09/97 (Figure 3 (d)).

Figure 1 (a) 11/29/96 Marine




Figure 1 (c) 01/13/97 Marine




Figure 1 (b) 12/06/96 Modified marine




Figure 1 (d) 01/09/97 Continental




Figure 1 (e) 02/07/97 Modified marine




Figure 1 (g) 03/13/97 Marine




Figure 1 (f) 01/31/97 Continental




Figure 1 (h) 03/19/97 Marine




Figure 1 (i) 04/03/97 Modified marine




Figure 1 (k) 05/08/97 Modified marine




Figure 1 (j) 04/24/97 Marine




Figure 1 (l) 05/02/97 Modified continental




Figure 2 (a) 11/29/96 Marine




Figure 2 (c) 01/13/97 Marine




Figure 2 (b) 12/06/96 Modified marine




Figure 2 (d) 01/09/97 Continental





Figure 2 (e) 02/07/97 Modified marine






Figure 2 (f) 01/31/97 Continental

Figure 2 (g) 03/13/97 Marine




Figure 2 (h) 03/19/97 Marine





Figure 2 (i) 04/03/97 Modified marine

Figure 2 (j) 04/24/97 Marine







Figure 2 (k) 05/08/97 Modified marine

Figure 2 (l) 05/02/97 Modified continental







Figure 3 (a) 11/29/96 Marine

Figure 3 (b) 12/06/96 Modified marine







Figure 3 (c) 01/13/97 Marine

Figure 3 (d) 01/09/97 Continental







Figure 3 (e) 02/07/97 Modified marine

Figure 3 (f) 01/31/97 Continental







Figure 3 (g) 03/13/97 Marine

Figure 3 (h) 03/19/97 Marine






Figure 3 (i) 04/03/97 Modified marine

Excitation @ 450 nm

1

Output

Output

Description

730

785

Figure 3 (j) 04/24/97 Marine




Figure 3 (k) 05/08/97 Modified marine

565

620

Wavelength (nm)

675

510





Figure 3 (1) 05/02/97 Modified continental






Figure 4 (a) 29 November 1996




Figure 4 (b) 06 December 1996




Figure 4 (c) 09 January 1997




Figure 4 (d) 13 January 1997




Figure 4 (e) 31 January 1997




Figure 4 (f) 07 February 1997




Figure 4 (g) 13 March 1997




Figure 4 (h) 19 March 1997




Figure 4 (i) 03 April 1997




Figure 4 (j) 24 April 1997




Figure 4 (k) 02 May 1997




Figure 4 (l) 08 May 1997